

# The role of binaries in the enrichment of the early Galactic halo

## II. Carbon-enhanced metal-poor stars – CEMP-no stars (*Corrigendum*)

T. T. Hansen<sup>1</sup>, J. Andersen<sup>2,3</sup>, B. Nordström<sup>2,3</sup>, T. C. Beers<sup>4</sup>, V. M. Placco<sup>4</sup>, J. Yoon<sup>4</sup>, and L. A. Buchhave<sup>5</sup>

<sup>1</sup> George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, and Department of Physics and Astronomy, Texas A&M University, College Station TX 77843, USA  
e-mail: [thansen@tamu.de](mailto:thansen@tamu.de)

<sup>2</sup> Dawn Cosmology Centre, The Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark  
e-mail: [ja@nbi.ku.dk](mailto:ja@nbi.ku.dk); [birgitta@nbi.ku.dk](mailto:birgitta@nbi.ku.dk)

<sup>3</sup> Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, 8000 Aarhus C, Denmark

<sup>4</sup> Department of Physics and JINA Center for the Evolution of the Elements, University of Notre Dame, Notre Dame, IN 46556, USA

<sup>5</sup> DTU Space, National Space Institute, Technical University of Denmark, Elektrovej 327, 2800 Lyngby, Denmark

A&A, 586, A160 (2016), <https://doi.org/10.1051/0004-6361/201527235>

**Key words.** Galaxy: formation – Galaxy: halo – stars: chemically peculiar – binaries: spectroscopic – ISM: structure – errata, addenda

In Hansen et al. (2016) the RV mean for HE 0020–1741 listed in Table 2 in the printed article is given incorrectly. The correct value is  $93.039 \text{ km s}^{-1}$ , in agreement with the individual RV values for the star listed in Table A.1.