The “memory” of the Oort cloud (Corrigendum)

Marc Fouchard¹, Arika Higuchi², Takashi Ito³, and Lucie Maquet¹

¹ LAL-IMCCE, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universités, UPMC Univ. Paris 06, Univ. Lille.,
 1 Impasse de l’Observatoire, 59000 Lille, France
 e-mail: marc.fouchard@univ-lille1.fr

² RISE Project Office, National Astronomical Observatory of Japan, 2-12 Hoshigaoka-cho, Mizusawa, Iwate 023-0816, Japan

³ CfCA, NAOJ, Mitaka, Tokyo 181-8588, Japan

A&A, 620, A45 (2018), <https://doi.org/10.1051/0004-6361/201833435>

Key words. comets: general – Oort Cloud – errata, addenda

The q axes of Figs. 1 and 3 are oriented downward, whereas they should be oriented upward. The figures themselves are correct, only the orientation of the axes is wrong.

Fig. 1. From top to bottom panels: distribution of comets in (a, q) , $(a, \cos(i_E))$, (a, ω_G) and (a, Ω_G) planes at four different epochs $t = 250$ Myr and 1, 2 and 4 Gyr. For visibility, the bin sizes are constant along any axis. However the density function is computed in number of comets per au for the perihelion distance and per au^{-1} for the semi-major axis. The data comes from the evolution of the 10 proto-Oort cloud.

Fig. 3. As Fig. 1 but for the D0 proto-Oort cloud.