LETTER TO THE EDITOR

The dry and carbon-poor inner disk of TW Hydrae: evidence for a massive icy dust trap

Arthur D. Bosman^{1,*} and Andrea Banzatti²

¹ Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands
e-mail: arbos@umich.edu

² Department of Physics, Texas State University, 749 N Comanche Street, San Marcos, TX 78666, USA

Received 5 September 2019 / Accepted 22 November 2019

ABSTRACT

Context. Gas giants accrete their envelopes from the gas and dust of proto-planetary disks, and therefore it is important to determine the composition of the inner few astronomical units, where most giant planets are expected to form.

Aims. We aim to constrain the elemental carbon and oxygen abundance in the inner disk ($R < 2.3$ AU) of TW Hya and compare with the outer disk ($R > 2.3$ AU) where carbon and oxygen appear underabundant by a factor of approximately 50.

Methods. Archival *Spitzer*-IRS and VLT-CRIRES observations of TW Hya were compared with a detailed thermo-chemical model, DALI. The inner disk gas mass and elemental C and O abundances were varied to fit the mid-infrared H_2 and H_2O line fluxes as well as the near-infrared CO line flux.

Results. Best-fitting models have an inner disk that has a gas mass of $2 \times 10^{-4} M_\odot$ with $C/H \approx 3 \times 10^{-6}$ and $O/H \approx 6 \times 10^{-6}$. The elemental oxygen and carbon abundances of the inner disk are about 50 times lower than in the interstellar medium and are consistent with those found in the outer disk.

Conclusions. The uniformly low volatile abundances imply that the inner disk is not enriched by ices on drifting bodies that evaporate. This indicates that drifting grains are stopped in a dust trap outside the water ice line. Such a dust trap would also form a cavity as seen in high-resolution submillimeter continuum observations. If CO is the major carbon carrier in the ices, dust needs to be trapped efficiently outside the CO ice line of ~ 20 AU. This would imply that the shallow submillimeter rings in the TW Hya disk outside of 20 AU correspond to very efficient dust traps. The most likely scenario is that more than 98% of the CO has been converted into less volatile species, for example CO_2 and CH_3OH . A giant planet forming in the inner disk would be accreting gas with low carbon and oxygen abundances as well as very little icy dust, potentially leading to a planet atmosphere with strongly substellar C/H and O/H ratios.

Key words. protoplanetary disks – line: formation – astrochemistry – stars: individual: TW Hya

1. Introduction

The elemental abundances of carbon and oxygen in proto-planetary disks are a vital input to planet formation models and combined with characterization of exoplanets can tell us about the formation history of those planets. The simple picture of elemental abundances changing statically at the ice lines of the main chemical species (e.g. Öberg et al. 2011) is being increasingly enriched by a number of modeling studies and observations. Chemical evolution can be efficient in changing the composition of the disk gas, changing the major carriers of carbon and oxygen (e.g., Eistrup et al. 2016; Schwarz et al. 2018; Bosman et al. 2018a). Furthermore, disk dynamics and dust evolution can efficiently transport the volatile component of the disk, changing the elemental composition of the gas and ice (Kama et al. 2016; Booth et al. 2017; Bosman et al. 2018a; Krijt et al. 2016, 2018).

As the interplay of the physical and chemical processes is complex, observations are needed to benchmark disk models and provide much needed input for planet formation models.

* *Present address:* Department of Astronomy, University of Michigan, 311 West Hall, 1085 S. University Avenue, Ann Arbor, MI 48109, USA.

Observations of CO and H_2O focusing on the outer regions of proto-planetary disks show that their abundances are up to two orders of magnitude lower than expected, implying that chemical and physical processes are indeed modifying the abundances of these species (Hogerheijde et al. 2011; Favre et al. 2013; Bergin et al. 2013; Miotello et al. 2017; Du et al. 2017). Nevertheless, planet formation models primarily need the composition around and within the H_2O ice line, as this is where we expect giant planets to form and accrete their atmospheres (e.g. Kennedy & Kenyon 2008; Cridland et al. 2017; Dawson & Johnson 2018).

Therefore, we aim to constrain the elemental abundances in the inner disk of TW Hya. The outer disk of TW Hya has been well studied and the elemental composition of the gas has been constrained from a variety of observations, including HD to trace the total disk mass, allowing for the measurement of absolute abundances (Hogerheijde et al. 2011; Bergin et al. 2013; Kama et al. 2016; Schwarz et al. 2016; Trapman et al. 2017). These efforts have shown that both the volatile carbon and oxygen abundance in the outer disk are lowered by a factor of about 50 compared to the interstellar medium (ISM). Observations of $^{13}C^{18}O$ in TW Hya show that while some CO comes from the

Table 1. H₂, CO, and H₂O fluxes from observations and modeling.

Molecular	Transition ⁽¹⁾	Wavelength (μ m)	Observed flux ⁽²⁾	Model flux ⁽²⁾	E_{up} (K)
H ₂	$J = 3-1$ ($S(1)$)	17.0	1.57 ± 0.05	1.7	1015
H ₂	$J = 4-2$ ($S(2)$)	12.3	1.21 ± 0.20	0.75	1681
CO	$\nu, J = 1, 10-0, 11$ ($\nu 1 P(10)$)	4.75	0.64 ± 0.05	0.72	3330
H ₂ O	$12_{5,8}-11_{2,9}$	17.12	<0.17	0.03	3243
H ₂ O	$11_{3,9}-10_{0,10}$	17.22	<0.29	0.05	2473
H ₂ O	$11_{2,9}-10_{1,10}$	17.36	<0.32	0.11	2396
H ₂ O	$9_{2,7}-8_{1,8}$	21.85	2.8 ± 0.2	6.7	1730
H ₂ O	$9_{4,5}-8_{3,6}$	28.23	2.1 ± 0.1	1.3	1929
	$9_{5,4}-8_{4,5}$				2122
H ₂ O	$8_{6,3}-7_{5,2}$	28.43	1.6 ± 0.2	1.2	2031
	$8_{6,2}-7_{5,3}$				2031
H ₂ O	$7_{7,0}-6_{6,1}$	28.6	1.7 ± 0.1	1.2	2006
	$7_{7,1}-6_{6,0}$				2006
H ₂ O	$6_{3,1}-5_{0,5}$	30.87	2.7 ± 0.2	1.8	934
	$8_{5,4}-7_{4,3}$				1806
H ₂ O	$6_{6,1}-5_{5,0}$	33	3.96 ± 0.46	4.0	1504
	$7_{5,2}-6_{4,3}$				1524

Notes. Observed CO line flux is taken from [Banzatti et al. \(2017\)](#), and H₂ and H₂O fluxes are extracted from the *Spitzer*-IRS spectra using the method in [Banzatti et al. \(2012\)](#). The table denotes the upper level energy of the dominant transitions of each blend according to our model. The H₂ S(2) line is contaminated by an OH line; the contribution of the OH line was estimated by taking the average flux of two adjacent OH lines. ⁽¹⁾In the case of H₂O, the dominant transition(s) of the feature. ⁽²⁾In units of 10^{-14} erg s⁻¹ cm⁻².

grains within the CO ice line, the CO abundance stays a factor of about 20 lower than the ISM abundance ([Zhang et al. 2017](#)), implying that a lot of the carbon is trapped on the grains or is converted into other species.

The physical and chemical structure of the inner disk of TW Hya has also been studied in detail. It is the disk that has been resolved at the highest physical resolution with ALMA ([Andrews et al. 2016](#)) and together with abundant photometry and infrared interferometry, there is a clear picture of the inner disk structure of TW Hya ([Andrews et al. 2012](#); [Menu et al. 2014](#); [Kama et al. 2016](#)). In the infrared, high signal-to-noise ratio *Spitzer*-IRS and VLT-CRIRES spectra have been taken. These observations include 3.2 km s^{-1} resolution observations of the CO $\nu = 1-0$ rovibrational band at $4.7 \mu\text{m}$, and detections of the CO₂ $15 \mu\text{m}$ Q -branch, pure rotational lines of H₂, H₂O, and OH, as well as a number of atomic hydrogen lines ([Najita et al. 2010](#); [Pontoppidan et al. 2008](#)). Modeling efforts have further constrained the inner disk gas mass using H₂ ([Gorti et al. 2011](#)) as well as the H₂O content of the inner disk ([Zhang et al. 2013](#)). These studies hint at abundances for CO and H₂O in the inner disk that are lower than expected from inner disk chemistry for gas of ISM composition, implying that elemental carbon and oxygen are depleted relative to the ISM.

Here we build on these studies, using the better-constrained outer disk structure and composition from [Trapman et al. \(2017\)](#), with an updated inner disk gap from ALMA observations ([Andrews et al. 2016](#)). Using this disk structure and the thermochemical code DALI we constrain the elemental carbon and oxygen abundance in the inner disk of TW Hya.

2. Methods

The lines and features, and their fluxes, that we consider for our modeling comparison are tabulated in Table 1. The CO flux for TW Hya is taken from [Banzatti et al. \(2017\)](#) based on VLT-CRIRES ([Kaeufl et al. 2004](#)) spectra presented in [Pontoppidan et al. \(2008, program ID 179.C-0151\)](#). The fluxes

for H₂ and H₂O are extracted from the *Spitzer*-IRS spectrum obtained from program GO 30300 using the method described in [Banzatti et al. \(2012\)](#). The spectrum has been published in [Najita et al. \(2010\)](#) and [Zhang et al. \(2013\)](#).

We start with the Dust and Lines (DALI, [Bruderer et al. 2012](#); [Bruderer 2013](#)) model of [Trapman et al. \(2017\)](#) for TW Hya (see also [Kama et al. 2016](#), model parameters are given in Table A.1). This model fits the SED, many far-infrared and submillimeter lines, and the ALMA ¹²CO $J = 3-2$ image. The most important lines are the HD 112 and $56 \mu\text{m}$ lines, constraining the outer disk mass, and many CO (isotopologue) lines constraining the carbon abundance. On top of these, some atomic carbon and oxygen fine-structure lines have also been fit, which constrain the elemental carbon and oxygen abundances in the outer disk at $\text{C}/\text{H} = 2.7 \times 10^{-6}$ and $\text{O}/\text{H} = 5.8 \times 10^{-6}$ which are a factor 50 lower than expected for the ISM. In this work we update the inner disk structure moving the outer edge of the gap from 4 to 2.4 AU in accordance with the bright submillimeter ring seen by [Andrews et al. \(2016\)](#) at this radius (assuming a distance of 54 parsec).

Figure 1 shows the surface density structure in the inner disk. The inner disk mass is varied by varying δ_{gas} in the inner disk using a model that has constant, low elemental abundances. The predicted H₂ lines for DALI are compared to the observed fluxes. The model that fits best is then used to constrain the elemental C and O abundances. We make the simplifying assumptions that the CO abundance in the inner disk scales linearly with the total elemental carbon abundance, that the H₂O abundance scales with the elemental oxygen that is not locked in CO, and that changing the CO and H₂O abundances does not significantly alter the gas temperatures. The elemental C and O abundances are varied by changing the CO and H₂O abundances by a factor δ_{C} and δ_{O} , respectively, above gas temperatures of T_{step} of 70, 150, and 500 K. Here, 70 and 150 K are the sublimation temperatures of CO₂ and H₂O respectively, whereas 500 K is a rough transition temperature for the release of carbon from a more refractory reservoir. The excitation is recalculated for these new

Fig. 1. Surface density structure in inner regions of the TW Hya model. δ_{gas} is the inner disk drop in gas surface density. The value $\delta_{\text{gas}} = 0.1$ used here best fits the H_2 observations. The red curves show the column averaged oxygen and carbon depletion factors in TW Hya. The red solid line shows a model with constant depletion. The dotted line shows the depletion profile assuming that carbon and oxygen return to the ISM values above $T_{\text{gas}} = 150$ K.

abundance structures and the CO $\nu 1$ P(10) line flux and the full H_2O spectrum between 12 and $34\ \mu\text{m}$ is extracted.

The excitation for H_2 is done in local thermodynamic equilibrium (LTE) as there are no collisional rate coefficients available for H_2 ; however, the low Einstein A coefficients of the H_2 lines mean that deviations from LTE are expected to be negligible (Wolniewicz et al. 1998). For H_2O and CO the local excitation–de-excitation balance is calculated explicitly. For CO, collision rate coefficients from Yang et al. (2010) for H_2 and Song et al. (2015) and Walker et al. (2015) for H are used (see also Bosman et al. 2019). For H_2O , the data file from the LAMDA database¹ (Schöier et al. 2005) is used.

3. Results

Figure 2 compares the emitting regions of the different species in the constant abundance model with $\delta_{\text{gas}} = 0.1$. There is a large overlap in the emitting areas, especially in the inner disk (< 2.4 AU). Water has the most confined emitting area and only probes the inner disk while CO and H_2 both also probe the outer disk, where the composition of the gas is already strongly constrained (Bergin et al. 2013; Kama et al. 2016; Trapman et al. 2017; Zhang et al. 2017).

Figure 3 compares the results of the DALI modeling with the observed fluxes. The H_2 fluxes constrain the inner disk gas mass to be around $1.7 \times 10^{-4} M_{\odot}$ ($\delta_{\text{gas}} \approx 0.1$), comparable to the value found in Gorti et al. (2011). The observations show a higher H_2 S(2)/S(1) line ratio compared to the models, indicating that the average gas temperature in the inner disk of the model is too low. The upper level energies of the CO (~ 3000 K) and H_2O (~ 1500 K) are similar to or higher than the upper level energies of the H_2 lines (~ 1000 and ~ 1700 K), therefore a higher temperature in the inner disk would lead to even lower inferred values for C/H and O/H.

Fig. 2. Map of the gas temperature in the inner region of the model together with the emitting areas of the H_2 S(1) line (black) the CO $\nu 1$ vibrational lines (orange) and the strongest line of the H_2O $33\ \mu\text{m}$ feature (purple). The green line shows $T_{\text{dust}} = 150$ K, the approximate location of the H_2O ice line. The emitting areas radially overlap in the inner disk.

Both the CO lines and H_2O lines fit well to an inner disk elemental abundance similar to the abundance found in the outer disk, which is a factor 50 lower compared to the ISM. Even a jump of a factor of two in the elemental carbon or oxygen abundance above 500 K can be ruled out based on the observed fluxes. The H_2O fluxes in Table 1 show that this model underpredicts the H_2O detections between 20 and $31\ \mu\text{m}$ by a factor of about two, which is in line with the gas temperature in the model being too low.

4. Discussion

4.1. Constraining the inner disk chemical structure

Here, using a more complete physical and chemical structure of both the inner and the outer disk, we can quantify the total H_2 mass, as well as the volatile carbon and oxygen abundance in the inner disk, confirming that the inner disk of TW Hya is both oxygen- and carbon-poor by a factor about 50 compared to the ISM. Furthermore, our modeling shows that there is no significant (factor 2 or more) increase in volatile carbon or oxygen in the inner disk. Thus, there is no sign of volatile release at the CO_2 or H_2O ice lines nor is there evidence of carbonaceous or silicate grain destruction at $T < 500$ K.

Both Gorti et al. (2011) and Zhang et al. (2013) studied the inner region of TW Hya using detailed modeling. They note that they overproduce the CO rovibrational lines by a factor of about two. Furthermore, Gorti et al. (2011) assume LTE excitation for CO, which generally underpredicts fluxes compared to models that include infrared pumping of the vibrational levels by the inner disk continuum emission (Bruderer et al. 2015; Bosman et al. 2017). These two factors together explain why our models need a CO or elemental carbon abundance that is almost two orders of magnitude lower than that of the ISM to fit the CO rovibrational line.

Zhang et al. (2013) use a low abundance in the inner disk $x_{\text{H}_2\text{O}} < 10^{-6}$ with a ring of high abundance H_2O at 4 AU (their ice line) to produce the *Spitzer* lines. Including such a ring and fitting the $33\ \mu\text{m}$ ($E_{\text{up}} = 1504$ K) feature would lead to inferring

¹ <http://strw.leidenuniv.nl/~moldata>

Fig. 3. Comparison of the H_2 line fluxes (*left*), CO rovibrational line flux (*middle*), and the H_2O $33\,\mu\text{m}$ feature flux (*right*) between models and data. *Left panel:* the amount of gas within 2.4 AU is varied. *Middle and right panels:* $\delta_{\text{gas}} = 0.1$ is used and C/H and O/H are varied respectively. Horizontal bands show the 1σ variation of the observed fluxes. The different coloured points denote different values for T_{step} . The temperature threshold above which the abundances are varied. The vertical lines in the *middle and right panel* show the volatile carbon and oxygen abundances assumed for the ISM (Meyer et al. 1998; Lacy et al. 2017).

lower oxygen abundances in the inner disk; however, it would also move the H_2O emission to colder gas, increasing the discrepancy between model and observations for the 22 and $28\,\mu\text{m}$ features. A constant-abundance model therefore fits the data better than a ring model.

4.2. Hiding C and O carriers?

The DALI models predict that CO and H_2O are the dominant gas-phase oxygen and carbon carriers in the inner disk. However, it is possible that carbon and oxygen are locked in other gaseous species. There are a few obvious candidates for hiding more carbon and oxygen in molecular gas: CO_2 , C_2H_2 , HCN, and CH_4 . There is a CO_2 feature detected in the *Spitzer* spectrum of TW Hya. Applying the model results from Bosman et al. (2017) to the detected feature retrieves a CO_2 abundance lower than 10^{-9} with respect to H_2 . The models in Bosman et al. (2017) do not have a gap, which if included would only increase the strength of the CO_2 lines. As such the CO_2 abundance $< 10^{-9}$ is a stringent upper limit and this molecule is not an abundant carrier of either carbon or oxygen.

C_2H_2 and HCN are detected in many proto-planetary disks (Salyk et al. 2011), however neither are detected in the spectrum of TW Hya. As the Q -branches of HCN and C_2H_2 around $14\,\mu\text{m}$ have similar upper level energies and Einstein A coefficients to the $15\,\mu\text{m}$ Q -branch of CO_2 , both HCN and C_2H_2 should have been detected if they are as abundant as CO_2 . The lack of observed features implies that both HCN and C_2H_2 contain less than 1% of the volatile carbon. The final possible gaseous carbon carrier is CH_4 . Observations of this molecule have so far only been successful in one disk, in which CH_4 lines have been detected in absorption (Gibb & Horne 2013). The CH_4 abundance is therefore not strongly constrained by observations. However, from chemical models CH_4 is not expected to be the most abundant carbon carrier in inner disk atmospheres (Walsh et al. 2015; Agúndez et al. 2018). As such, with CO and H_2O we trace the bulk of gaseous carbon and oxygen in the inner disk.

4.3. Implications of uniform depletion

The low elemental abundances in the inner disk imply one of two things: either the entire disk is depleted in volatiles, implying

dust grains with almost no ices, or volatiles are efficiently locked in ices on solids that are not transported through the disk.

To deplete the entire disk in volatiles, a period of very strong radial drift coupled with strong radial mixing is necessary (e.g., Booth et al. 2017). In such a scenario, the icy grains drift from the mass reservoir in the outer disk into the regions of the disk where the ices desorb. This leads to enhancements in C/H and O/H in the inner regions of the disk. After a few million years all of this high-C/H and high-O/H gas is accreted onto the star and the disk is left depleted in carbon and oxygen. However, a disk like this will also be strongly depleted in solid material as the grains that transport the ices inward will continue to drift into the star leading to dust depletions at least as high as the volatile element depletion. This scenario is therefore very unlikely for TW Hya, which has a massive dust component and a gas-to-dust ratio of around 100 (Bergin et al. 2013).

As efficient transport is excluded, this leaves 98% of the carbon and oxygen locked in solids that are not efficiently transported. Assuming that CO and H_2O are the dominant oxygen and carbon carriers in the outer disk, it is necessary to stop CO from being transported over the CO ice line by a dust trap at a location larger than where the CO ice line is located. For TW Hya this would imply that one of the shallow millimeter continuum rings of TW Hya outside of 20 AU corresponds to an efficient dust trap. An efficient dust trap at > 20 AU should result in a disk with a large cavity. As dust is abundant down to 2.4 AU, a second dust trap would be necessary at that location, which would trap H_2O -rich grains.

In this scenario H_2O would be far more efficiently locked in large grains than CO as the former is frozen out in a larger fraction of the disk. However, if gaseous CO is efficiently converted into less volatile species, especially CO_2 and CH_3OH , then the CO- and H_2O -depletion fractions are likely to be more similar and an efficient dust trap around 20 AU is no longer strictly necessary. The age of TW Hya of 10 Mr, is long enough to convert large amounts of CO into other species (Donaldson et al. 2016; Bosman et al. 2018b). In this case, the dust should not be allowed to pass the CO_2 or CH_3OH ice lines. These ice lines are at nearly the same location as the H_2O ice line at the inner edge of the outer disk. As such, a dust trap at the innermost submillimeter ring would trap all of the icy CO_2 , CH_3OH , and H_2O in the outer disk. A schematic representation of this is given in Fig. 4.

Fig. 4. Schematic of the TW Hya disk showing the locations that should trap the oxygen- and carbon-bearing molecules. Oxygen needs to be trapped outside of 2.4 AU and is most likely trapped in the dust ring at that location. Carbon can either be trapped at the same location, with CO₂ being the most probable carrier, or at larger radii outside the CO ice line, with CO being the most probable carrier.

In summary, we find that the elemental carbon and oxygen abundances in the inner disk are lower by a factor of approximately 50 compared to the ISM. Even at temperatures of 500 K, the gaseous carbon or oxygen elemental abundances cannot have increased by a factor of two, strongly constraining the release of volatiles and grain destruction up to 500 K. This is interpreted as the dust trap responsible for the dust-free cavity also trapping the major carbon- and oxygen-bearing ices outside of 2.4 AU. A planet currently accreting gas in the gap will accrete very low amounts of carbon and oxygen, while possibly accreting ISM concentrations of nitrogen and noble gasses. Depending on the accretion history this planet could have a substellar C/H and O/H abundance.

Acknowledgements. Astrochemistry in Leiden is supported by the Netherlands Research School for Astronomy (NOVA). This work is partly based on observations made with CRIRES on ESO telescopes at the Paranal Observatory under program ID 179.C-0151. This work is based in part on observations made with the *Spitzer* Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. This project has made use of the SciPy stack (Virtanen et al. 2019), including NumPy (Oliphant 2006) and Matplotlib (Hunter 2007).

References

- Agúndez, M., Roueff, E., Le Petit, F., & Le Bourlot, J. 2018, *A&A*, **616**, A19
Andrews, S. M., Wilner, D. J., Hughes, A. M., et al. 2012, *ApJ*, **744**, 162
Andrews, S. M., Wilner, D. J., Zhu, Z., et al. 2016, *ApJ*, **820**, L40
Banzatti, A., Meyer, M. R., Bruderer, S., et al. 2012, *ApJ*, **745**, 90
Banzatti, A., Pontoppidan, K. M., Salyk, C., et al. 2017, *ApJ*, **834**, 152
Bergin, E. A., Cleeves, L. I., Gorti, U., et al. 2013, *Nature*, **493**, 644
Booth, R. A., Clarke, C. J., Madhusudhan, N., & Ilee, J. D. 2017, *MNRAS*, **469**, 3994
Bosman, A. D., Bruderer, S., & van Dishoeck, E. F. 2017, *A&A*, **601**, A36
Bosman, A. D., Tielens, A. G. G. M., & van Dishoeck, E. F. 2018a, *A&A*, **611**, A80
Bosman, A. D., Walsh, C., & van Dishoeck, E. F. 2018b, *A&A*, **618**, A182
Bosman, A. D., Banzatti, A., Bruderer, S., et al. 2019, *A&A*, **631**, A133
Bruderer, S. 2013, *A&A*, **559**, A46
Bruderer, S., van Dishoeck, E. F., Doty, S. D., & Herczeg, G. J. 2012, *A&A*, **541**, A91
Bruderer, S., Harsono, D., & van Dishoeck, E. F. 2015, *A&A*, **575**, A94
Cridland, A. J., Pudritz, R. E., Birnstiel, T., Cleeves, L. I., & Bergin, E. A. 2017, *MNRAS*, **469**, 3910
Dawson, R. I., & Johnson, J. A. 2018, *ARA&A*, **56**, 175
Donaldson, J. K., Weinberger, A. J., Gagné, J., et al. 2016, *ApJ*, **833**, 95
Du, F., Bergin, E. A., Hogerheijde, M., et al. 2017, *ApJ*, **842**, 98

- Eistrup, C., Walsh, C., & van Dishoeck, E. F. 2016, *A&A*, **595**, A83
Favre, C., Cleeves, L. I., Bergin, E. A., Qi, C., & Blake, G. A. 2013, *ApJ*, **776**, L38
Gibb, E. L., & Horne, D. 2013, *ApJ*, **776**, L28
Gorti, U., Hollenbach, D., Najita, J., & Pascucci, I. 2011, *ApJ*, **735**, 90
Hogerheijde, M. R., Bergin, E. A., Brinch, C., et al. 2011, *Science*, **334**, 338
Hunter, J. D. 2007, *Comput. Sci. Eng.*, **9**, 90
Kaeufl, H. U., Ballester, P., Biereichel, P., et al. 2004, in *Ground-based Instrumentation for Astronomy*, eds. A. F. M. Moorwood, & M. Iye, *Proc. SPIE*, **5492**, 1218
Kama, M., Bruderer, S., van Dishoeck, E. F., et al. 2016, *A&A*, **592**, A83
Kennedy, G. M., & Kenyon, S. J. 2008, *ApJ*, **673**, 502
Krijt, S., Ciesla, F. J., & Bergin, E. A. 2016, *ApJ*, **833**, 285
Krijt, S., Schwarz, K. R., Bergin, E. A., & Ciesla, F. J. 2018, *ApJ*, **864**, 78
Lacy, J. H., Sneden, C., Kim, H., & Jaffe, D. T. 2017, *ApJ*, **838**, 66
Menu, J., van Boekel, R., Henning, T., et al. 2014, *A&A*, **564**, A93
Meyer, D. M., Jura, M., & Cardelli, J. A. 1998, *ApJ*, **493**, 222
Miotello, A., van Dishoeck, E. F., Williams, J. P., et al. 2017, *A&A*, **599**, A113
Najita, J. R., Carr, J. S., Strom, S. E., et al. 2010, *ApJ*, **712**, 274
Öberg, K. I., Boogert, A. C. A., Pontoppidan, K. M., et al. 2011, *ApJ*, **740**, 109
Oliphant, T. 2006, *NumPy: A Guide to NumPy* (USA: Trelgol Publishing)
Pontoppidan, K. M., Blake, G. A., van Dishoeck, E. F., et al. 2008, *ApJ*, **684**, 1323
Salyk, C., Pontoppidan, K. M., Blake, G. A., Najita, J. R., & Carr, J. S. 2011, *ApJ*, **731**, 130
Schöier, F. L., van der Tak, F. F. S., van Dishoeck, E. F., & Black, J. H. 2005, *A&A*, **432**, 369
Schwarz, K. R., Bergin, E. A., Cleeves, L. I., et al. 2016, *ApJ*, **823**, 91
Schwarz, K. R., Bergin, E. A., Cleeves, L. I., et al. 2018, *ApJ*, **856**, 85
Song, L., Balakrishnan, N., Walker, K. M., et al. 2015, *ApJ*, **813**, 96
Trapman, L., Miotello, A., Kama, M., van Dishoeck, E. F., & Bruderer, S. 2017, *A&A*, **605**, A69
Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2019, *ArXiv e-prints* [arXiv:1907.10121]
Walker, K. M., Song, L., Yang, B. H., et al. 2015, *ApJ*, **811**, 27
Walsh, C., Nomura, H., & van Dishoeck, E. 2015, *A&A*, **582**, A88
Wolniewicz, L., Simbotin, I., & Dalgarno, A. 1998, *ApJS*, **115**, 293
Yang, B., Stancil, P. C., Balakrishnan, N., & Forrey, R. C. 2010, *ApJ*, **718**, 1062
Zhang, K., Pontoppidan, K. M., Salyk, C., & Blake, G. A. 2013, *ApJ*, **766**, 82
Zhang, K., Bergin, E. A., Blake, G. A., Cleeves, L. I., & Schwarz, K. R. 2017, *Nat. Astron.*, **1**, 0130

Appendix A: DALI model

Table A.1. Parameters of the TW Hya model.

Stellar parameters	
Stellar mass	$0.74 M_{\odot}$
Stellar luminosity	$1 L_{\odot}$
X-ray luminosity	$10^{30} \text{ erg s}^{-1}$
Disk parameters	
Disk mass	$0.025 M_{\odot}$
Critical radius (R_c)	35 AU
Surface density slope (γ)	1
Scale height at R_c (h_c)	0.1 rad
Flaring angle (ψ)	0.3
Inner radius	0.05 AU
Gap inner radius	0.3 AU
Gap outer radius	2.4 AU
Inner disk gas-to-dust	100
Inner disk gas depletion	$[10^{-5}-1]$
Small dust size	$0.005-1 \mu\text{m}$
Small dust fraction	0.01
Large dust size	$0.005-1000 \mu\text{m}$
Large dust fraction	0.99
Large dust settling factor	0.2