Small glitches and other rotational irregularities of the Vela pulsar (Corrigendum)

C. M. Espinoza¹, D. Antonopoulou^{2,3}, R. Dodson⁴, M. Stepanova¹, and A. Scherer⁵

¹ Departamento de Física, Universidad de Santiago de Chile (USACH), Estación Central, Santiago 9170124, Chile
e-mail: crisobal.espinoza.r@usach.cl

² Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, ul. Bartycka 18, 00-716 Warsaw, Poland

³ Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester, Manchester M13 9PL, UK

⁴ International Centre for Radio Astronomy Research, University of Western Australia, Crawley, WA 6009, Australia

⁵ Instituto de Astrofísica, Pontificia Universidad Católica de Chile, 782-0436 Macul, Santiago, Chile

A&A 647, A25 (2021), <https://doi.org/10.1051/0004-6361/202039044>

Key words. stars: neutron – pulsars: general – pulsars: individual: PSR B0833–45 – stars: rotation – errata, addenda

There was an error with the plotted lines that indicate detection capabilities in quadrants I and III of Fig. 9 (for events with $\Delta\nu$ and $\Delta\dot{\nu}$ of equal signs). The correct figure and updated caption is below.

Section 6: To accommodate the correction to Fig. 9, the last sentence of the second paragraph must read “For the other events (quadrants I and III) we used the condition $\phi_g(t = 10 \text{ d}) > \sigma_{\phi}^{\text{max}}$ as an indication for event sizes that would be under typical noise levels”.

Fig. 9. Sizes of all the irregularities selected for all possible combinations of signs for $\Delta\nu$ and $\Delta\dot{\nu}$, including cases in which one of the parameters was not detected ($\Delta\nu = 0$ or $\Delta\dot{\nu} = 0$, which are plotted as a letter x and crosses, respectively). These results are for $N_d = 10$ d, and the known Vela glitches in the dataset were measured as for Fig. 2. The straight lines represent detection limits (events in the shaded areas are unlikely to be detected) when considering average noise levels. The curved dotted lines give an indication of the event sizes below which deviations to the timing model will remain under typical noise levels for 10 d since the event epoch (see Sect. 6).