

PRESS RELEASE

Released on January 15, 2009

First ground-based detection of light from transiting exoplanets

Based on the articles "Ground-based K-band detection of thermal emission from the exoplanet TrES-3b", by E.J.W. De Mooij and I.A.G. Snellen, and "Ground-based secondary eclipse detection of the very-hot Jupiter OGLE-TR-56b", by D. K. Sing and M. Lopez-Morales

Published in *Astronomy & Astrophysics Letters*, 2009, volume 493-2, pp. L31-L38.

This week, *Astronomy & Astrophysics* is publishing the first ground-based detections of thermal emission from transiting exoplanets. Up to now, detections of this kind have only been made from space. Two independent teams are now presenting detections with ground-based telescopes of the thermal emission from transiting hot Jupiters.

Transiting exoplanets are routinely detected when they pass in front of their parent star as viewed from the Earth, which only happens by chance. The transit event causes a small drop in the observed starlight, which can then be detected. Fifty-five exoplanets have been detected this way since the observation of the first transiting planet HD 209458 b in 1999. When the planet revolves around its star or when it goes behind, the light coming from the system also varies, though the resulting smaller modulation is much harder to detect. This is mostly due to the small amount of light emitted by these exoplanets which are believed to be as dark as coal and reflect little of the incoming starlight. Fortunately, some of these planets are very hot, thus emitting light, mostly at infrared wavelengths. Up to now, detections of this kind have only been made using the Spitzer infrared space telescope. This week, *Astronomy & Astrophysics* is publishing the two first ground-based detections of thermal emission from transiting, hot-Jupiter exoplanets, from two independent teams of astronomers that used different approaches.

One team includes Ernst De Mooij and Ignas Snellen (University of Leiden, Netherlands) who used the William Herschel 4.2 meter telescope in La Palma (Canary Islands, Spain) to observe the star TrES-3 and its planet TrES-3b. To be able to detect the light coming from the planet, they observed the planet exactly at the time when it passes behind the star. They observed the event at infrared wavelengths, where the planet is at its brightest compared to the star (even if the planet is still much fainter than the star!) As they detected the light coming from the planet, they estimated the temperature of its atmosphere to about 2000 Kelvins. This indicates that the day side of the planet is extremely hot.

The other team, involving David Sing (IAP, France) and Mercedes Lopez-Morales (Carnegie Institution of Washington, USA), had a different approach. They looked at a much fainter star and its planet, OGLE-TR-56b. This planet is one of the most irradiated planets known so far, both because the planet is very close to the star and because the star is very hot. To detect the slight modulation in light that occurs when the planet passes behind its star, they used the 8 meter Very Large Telescope (ESO, Chile) and the 6.5 meter Magellan Telescopes (Las Campanas, Chile) and were able to observe this event at visible wavelengths. Indeed, the planet OGLE-TR-56b is heated so much by its star that it emits detectable amounts of

light in the visible wavelengths, and not only in the infrared as TrES-3b does. Hence, Sing and Lopez-Morales measured the record-high temperature of a planetary atmosphere: 2700 Kelvins. As in the case of TrES-3b, such a high day-side temperature indicates that winds cannot redistribute the heat efficiently from the day side to the night side.




Fig. 1. Artist's impression of the star OGLE-TR-56 and its planet, as the planet is passing behind the star. © D. Sing (IAP) / A&A.

These two independent results are very interesting for astronomers and planetary scientists because they allow a direct probe of the temperature of these planetary atmospheres, and because they show that such measurements can be made from ground-based observatories, and not only when using space telescopes.

Ground-based K-band detection of thermal emission from the exoplanet TrES-3b, by E.J.W. De Mooij and I.A.G. Snellen. Published in *Astronomy & Astrophysics Letters*, 2009, volume 493-2, pp. L31-L34

Ground-based secondary eclipse detection of the very-hot Jupiter OGLE-TR-56b, by D. K. Sing and M. Lopez-Morales. Published in *Astronomy & Astrophysics Letters*, 2009, volume 493-2, pp. L35-L38

Free version of the articles in PDF format: [De Mooij & Snellen](#), [Sing & Lopez-Morales](#)
Electronic edition of the press release and PDF files available at <http://www.aanda.org>

Contacts:

- Science:

Ernst de Mooij
Leiden University, Netherlands
Email: demooij (at) strw.leidenuniv.nl
Phone: +31 (0)71 527 8492

Dr. Mercedes Lopez-Morales
Carnegie Institution of Washington, USA
Email: mercedes (at) dtm.ciw.edu
Phone: +34 922 605 379

Dr. David. Sing
Institut d'Astrophysique de Paris, CNRS/UPMC
France
Email: sing (at) iap.fr
Phone: +33 (0)1 44 32 80 22

Dr. Ignas Snellen
Leiden University, Netherlands
Email: snellen (at) strw.leidenuniv.nl
Phone: +31 (0)71 527 5838

- Press office:

Dr. Jennifer Martin
Astronomy & Astrophysics
61, avenue de l'Observatoire
75014 Paris, France

Email: aanda.paris (at) obspm.fr
Phone: +33 (0)1 43 29 05 41

To subscribe to the *Astronomy & Astrophysics* press release mailing list, please contact Jennifer Martin at [aanda.paris \(at\) obspm.fr](mailto:aanda.paris@obspm.fr).